



Impact of Self-Learning Nutrition Education on Improving Awareness and Knowledge of Age-Related Macular Degeneration, and the Role of Lutein/Zeaxanthin in Eye Health

Kansuda Wunjuntuk, PhD (Corresponding author)

Faculty of Agriculture, Kasetsart University

E-mail: fagrkdw@ku.ac.th

Wipada Saelao, Graduate Student

Faculty of Agriculture, Kasetsart University

E-mail: wipada.sae@ku.th

Assistant Professor Sutithep Siripipattanakul, PhD

Faculty of Education, Kasetsart University

E-mail: fedustt@ku.ac.th

Associate Professor Prapaisri Sirichakwal, PhD

Institute of Nutrition, Mahidol University

E-mail: prapaisri.sir@mahidol.ac.th

Associate Professor Prut Hanutsaha, MD

Faculty of Medicine Ramathibodi Hospital, Mahidol University

E-mail: prut.han@mahidol.ac.th

Abstract

Age-related macular degeneration (AMD) is a condition affecting older people and involves the irreversible loss of central vision. Studies shown that increased dietary intake of lutein/zeaxanthin (L/Z) is associated with a lower risk of AMD. The purposes of this study were to 1) develop a set of 2 self-learning nutrition education consists of an AMD video and a role of L/Z in eye health video. 2) study satisfaction with the set of 2 videos in adults and the elderly 3) Compare awareness and knowledge of AMD, and the role of L/Z in eye health before and after viewing each video. This study was conducted in a quasi-experimental design, setting in Bangkok, Thailand. A voluntary sample of 103



adults and the elderly aged between 50 and 75 was recruited during April – May 2018. Participants were asked to complete a pre-test which was a questionnaire for awareness and knowledge of AMD prior to viewing the AMD video, and to complete a pre-test questionnaire for the role of L/Z in eye health prior to viewing the role of L/Z in eye health video. After self-learning through each of video, participants were asked to complete a post-test which was the same questionnaire as the pre-test and completed a satisfaction questionnaire for that video. The research instruments have been verified by 3 experts, having IOC value of 0.91. The reliability of instrument was tested using Cronbach's alpha coefficient, $\alpha = 0.879$. The data was analyzed by mean, standard deviation and t-test. Results revealed that 1) Participants had an overall satisfactory score on both of AMD video and the role of L/Z in eye health video at extremely satisfied level ($\bar{x}=4.60$, S.D.=0.427 and $\bar{x}=4.68$, S.D.=0.429, respectively). 2) After the self-learning through the AMD video, participants showed a significantly higher score in their awareness and knowledge of AMD compared to their score before self-learning ($\bar{x}=11.63$, S.D.=2.347 and $\bar{x}=3.78$, S.D.=2.642, respectively) at $p<0.05$ ($t= 25.148$, P-value= 0.000). 3) After the self-learning through the role of L/Z in eye health video, participants also showed a significantly higher score in their awareness and knowledge of the role of L/Z in eye health compared to their score before self-learning ($\bar{x}=12.88$, S.D.=9.474 and $\bar{x}=5.10$, S.D.=3.661, respectively) at $p<0.05$ ($t= 8.190$, P-value= 0.000). This can be concluded that the developed self-learning nutrition education were readily accepted by the adults and elderly. In addition, the nutrition intervention has shown a positive impact on improving awareness and knowledge of AMD, and the role of L/Z in eye health for adults and the elderly.

Keywords: Nutrition Education, Lutein/Zeaxanthin, Adults and the Elderly, Age-Related Macular Degeneration.

1. Introduction

Age-related macular degeneration (AMD) is a condition affecting older people and involves the loss of the person's central field of vision (World Health Organization [WHO], 2019). According to WHO (2019), AMD ranks third as a cause of blindness after cataracts and glaucoma, and it is the primary cause of blindness in industrialized countries. Regarding the 4th National Survey of Blindness, Low Vision and Major Eye Diseases in Thailand (2007), AMD also marked as the 3rd leading cause of blindness after cataracts and glaucoma, with approximately 799,296 Thai people suffering from AMD and 21,425 people having had irreversible blindness associated with AMD (Yenjit et al., 2007). The incidence of AMD is expected to rise with the aging of the population. While anti-vascular endothelial growth factor (anti-VEGF) therapy has been introduced for the wet form of the disease, a therapeutic regimen for the dry form, which affects most AMD patients, is still lacking. Therefore, preventive strategies such as education to



www.mecsj.com

raise awareness and knowledge about AMD, emphasis on lifestyle changes that can impact the elderly's eye health are becoming prudent for policy makers and health care practitioners.

Lifestyle practices such as eating a healthy diet high in green leafy vegetables, eating fish high in omega-3 fatty acids, avoid smoking, maintaining normal blood pressure, cholesterol level as well as exercise regularly, can play a significant role in reducing the risk of developing AMD (National Eye institute, 2019). The two carotenoids, namely lutein and zeaxanthin (L/Z), which are antioxidants and only 2 carotenoids located in the eye retina, may protect against AMD (American Optometric Association [AOA], 2019). Numerous studies have identified L/Z to be essential components for eye health. One of the first large studies on carotenoids is the Eye Disease Case Control Study, in which diets were compared as the risk factors for developing AMD. Results revealed a significantly lower risk for developing AMD in people with high amounts of L/Z in their blood. Also, those people eating a diet with the most L/Z (as much as 5.8 milligrams (mg) per day) had a significantly lower risk for AMD than those whose diet contained the least amount (as low as 1.2 mg per day) (Seddon et al., 1994). In addition, dietary studies have confirmed the association between frequent consumption of spinach or collard greens, particularly good sources of L/Z, and a lower risk of AMD. An analysis of a national dietary study called the 3rd National Health and Nutrition Examination Survey (NHANES III) in the US showed that consuming 6 mg per day of dietary L/Z was associated with a reduced risk for developing AMD (Marse-Perlman et al., 2001; AOA, 2019)

L/Z are the most common carotenoids found in green leafy vegetables (e.g. kale, spinach, broccoli, peas and lettuce), egg yolks and corn (Abdel-Aal et al., 2013). Given the positive association between L/Z and AMD, it seems prudent for people to obtain higher amounts of these nutrients from their daily diet especially from green leafy vegetables. However, majority of populations especially Thai people are not consuming enough fruits and vegetables each day. According to the 5th Thai National Health Examination Survey (NHES V), almost half of the population are consuming less than 2 servings of vegetables each day. Furthermore, older adults tend to consume fewer vegetables than younger adults (HSRI, 2019). Although there is no recommended daily intake for L/Z for Thai people, the studies referenced here suggest an intake of 6 mg or more per day of dietary L/Z to decrease the risk of developing AMD (AOA, 2019).

Nutritional education is a process by which people are assisted in making decisions about eating. Its ultimate goals are to improve the recipients' nutritional status or other aspects of their nutritional well-being through both knowledge acquisition, and behavior change skills (FAO, 2019). According to Parker et al. (2011), the video educational delivery strategy was mostly preferred by older adults, especially those with limited income and they also reported that the video was more enjoyable and found the educational content more stimulating when compared to PowerPoint and handout educational delivery strategy. Therefore, researchers were interested in developing a set of nutrition education videos consisting of the AMD disease video and the role of L/Z nutrients in eye health




Figure (2): Pictures show how the sight of someone with AMD can be affected. (ForeseeHome, 2019; Talley Eye Institute, 2019)



Figure (3): The ophthalmologist demonstrated on how to detect AMD by one self at home. (Thammasat Eye Center, 2019)



Figure (4): AMD risk factors and lifestyle practice that can prevent AMD. (OneBite Wellness, 2019; Macula Center, 2019)

2.1.1.2 The role of L/Z in eye health video (Figure 5-13) is a 20-minute self-learning video. There was a modulator interviewed a nutritionist, discussing about what people should know about the role of nutrients in eye health, what L/Z is, why L/Z can prevent AMD, the recommended daily intake for L/Z, the dietary sources and suggested meal plans to get enough dietary L/Z. Researchers referred the amount of dietary L/Z found in foods which demonstrated in the videos from the USDA Food Composition Databases (2017). However, for Thai dishes, we referred to results of Busaba & Siriporn (2012). To make the video to be easy recognized for L/Z-rich foods, researchers divided L/Z-rich foods into 5 groups including green leafy vegetables, legumes and nuts, orange-red colored vegetables, grains, and egg. Amount of L/Z, percentage of the suggested daily intake for L/Z and rating for each item of L/Z-rich food were described in table 1.



Table (1): Amount of dietary L/Z (mg.) found in each item of L/Z-rich foods which were categorized into 5 groups.

L/Z rich foods	Food Quantity (edible portion)	Amount of L/Z (mg)	% Suggested daily intake for L/Z*	Rating (number of stars**)
Group I: Green Leafy Vegetables				
Cooked Kale	1 ladle	7.6	130	6
Cooked Chinese Spinach	1 ladle	5.0	80	5
Cos (lettuce)	2 cups	1.5	25	1.5
Cooked Broccoli	2 ladles	1.2	20	1
Cooked Asparagus	2 ladles	0.9	15	1
Fresh Basil	20 leaves	0.6	10	0.5
Group II: Legumes and Nuts				
Cooked Edamame Beans	1 cup	2.5	40	2.5
Cooked Green Peas	½ cup	2.2	35	2
Cooked Green Beans	2 ladles	0.8	15	1
Pistachio	½ cup	0.7	10	0.5
Group III: Orange-red colored vegetables				
Cooked Pumpkin	2 ladles	1.1	20	1
Fresh Chili	1 tbsp	0.8	15	1
Cooked Carrot	2 ladles	0.6	10	0.5
Chili Powder	1 tsp	0.4	6	0.5
Group IV: Grains				
Cooked Sweet Corn	2 ladles	1.1	20	1
Group V: Egg				
Boiled Egg	2 eggs	0.4	6	0.5

*percentage of the suggested daily intake for L/Z (AOA, 2019)

**Rating (number of stars) for each item of L/Z-rich foods (consume totally 6 stars/day, or 100% of suggested daily intake for L/Z will obtain adequate amount of L/Z per day)



www.mecsj.com



Figure (5): The modulator interviewed the nutritionist, discussing about the role of nutrients in eye health.



Figure (6): Eating 2 ladles of cooked Chinese kale or Chinese spinach, may help to achieve 100% of dietary L/Z recommended per day.



Figure (7): 2 ladles of cooked broccoli or 2 cups of lettuce provide about 20-25% of dietary L/Z recommended per day.



Figure (8): 1 tablespoon of chili provide about 15% of dietary L/Z recommended per day.



Figure (9): 2 ladles of cooked corn or pumpkin provide about 20% of dietary L/Z recommended per day.



Figure (10): 2 ladles of cooked carrot provide about 10% of dietary L/Z recommended per day.



Figure (11): 1/2 cup of cooked green peas provide about 35% of dietary L/Z recommended per day.



Figure (12): A cup of cooked edamame beans provide about 40% of dietary L/Z recommended per day.



Figure (13): A boiled egg provides about 3-5% of dietary L/Z recommended per day.

2.1.2 A questionnaire to collect personal general information of participants, including gender, age, marriage status, educational background, occupation, health status and family history in relation to AMD.

2.1.3. A questionnaire to access satisfaction with the set of two self-learning nutrition education videos.



2.1.4. A questionnaire to assess awareness and knowledge of AMD, and a questionnaire to assess awareness and knowledge of the role of L/Z in eye health.

The questionnaire for awareness and knowledge of AMD comprised 15 questions, while the questionnaire for awareness and knowledge of the role of L/Z in eye health comprised 16 questions. All questionnaires were verified by 3 experts (an ophthalmologist, a nutritionist and an expert in elderly education media) having an IOC value of 0.91. The reliability of the instrument was tested on 30 adults and elderly in test group using Cronbach's alpha coefficient, $\alpha = 0.879$. The data was analyzed by mean, standard deviation and paired sample t-test.

2.2 Human research protection

This study was approved by the Kasetsart University research ethics committee, carried out in compliance with the international guidelines for human research protection as a declaration of Helsinki, the Belmont Report, CIOMS guideline, and international conference on harmonization in good clinical practice (ICH-GCP). The study code is KUREC-HS61/007. The approval number is COA61/029, with an approval expiry date of 1st April 2019. Participants in this study were counseled, explained the purpose of the study, signed the informed consent, and have the right to terminate the study whenever required without any impact. The study was conducted only on voluntary adults and elderly.

2.3 Data collection

2.3.1 The participant was asked to fill out the questionnaire about their personal information, filled out the pre-tests which were a questionnaire about awareness and knowledge of AMD, and a questionnaire about the role of L/Z in eye health prior to the self-learning.

2.3.2 Participants viewed the AMD video, then answered the post-test which was the same set of questions as pre-test and filled out a questionnaire for satisfaction towards this self-learning video.

2.3.3 Participants viewed the role of L/Z in eye health video, then answered the post-test which was the same set of questions as pre-test and filled out a questionnaire for satisfaction towards this self-learning video.



2.4 Data analysis

Statistical significance was determined to be a level of .05, and analyzed the following data:

2.4.1 Personal information: data analyzed by descriptive statistics, mean and standard deviation.

2.4.2 Satisfaction towards the set of two self-learning videos, data analyzed by descriptive statistics, mean and standard deviation.

2.4.3 Comparison of the mean scores for awareness and knowledge of AMD, and the role of L/Z in eye health, between the score before and after viewing the self-learning videos, data analyzed by paired-sample t-test.

3. Results

A total of 83% of the participants were female and 17% were male, 73% of participants were aged between 50-60 years, while 27% were aged between 61-75 years with an average age of 57 years. A total of 60% of participants are married, 65% of participants were under graduated. In addition, 12% of the participants reported on having a family member with AMD. After viewing the self-learning nutrition education videos, the participants found satisfaction in various aspects as described in table 2 & 3 at an extremely satisfied level. The results for satisfaction in various aspects were in line with the results for overall satisfaction. Participants also found an overall satisfaction towards the AMD video and the L/Z role in eye health video at an extremely satisfied level ($\bar{x}=4.60$, S.D.=0.427 and $\bar{x}=4.68$, S.D.=0.429, respectively). (Table 2, 3)

Table (2): Satisfaction with various aspects of the AMD videos of the participants after viewing the self-learning video.

Satisfaction with the AMD video	Mean	S.D.	Satisfaction level
1. Easy understanding of the story	4.59	0.550	Extremely satisfied
2. Content matched to topic	4.61	0.528	Extremely satisfied
3. Clear illustrations matched to content	4.63	0.524	Extremely satisfied
4. Illustration helped to easily understand content	4.62	0.562	Extremely satisfied
5. Suitable sound	4.61	0.660	Extremely satisfied
6. Suitable and easy understandable language	4.73	0.469	Extremely satisfied



Table (2): Cont.

Satisfaction with the AMD video	Mean	S.D.	Satisfaction level
7. Suitable and clear easy reading font and font size	4.50	0.715	Extremely satisfied
8. Interesting story that enhanced engagement	4.53	0.607	Extremely satisfied
9. Suitable video length	4.53	0.623	Extremely satisfied
10. Video encouraged further learning	4.54	0.623	Extremely satisfied
11. The content was practical in daily life	4.66	0.552	Extremely satisfied
12. Overall satisfaction	4.60	0.427	Extremely satisfied

Satisfaction score: 5=extremely satisfied, 4=very satisfied, 3=moderately satisfied, 2=slightly satisfied, 1=not at all

Table (3): Satisfaction with various aspects of the role of L/Z in eye health videos of participants after viewing the self-learning video

Satisfaction with the L/Z video	Mean	S.D.	Satisfaction level
1. Easy understanding of the story	4.66	0.552	Extremely satisfied
2. Content matched to topic	4.70	0.461	Extremely satisfied
3. Clear illustrations matched to content	4.71	0.478	Extremely satisfied
4. Illustrations helped to easily understand content	4.75	0.437	Extremely satisfied
5. Suitable sound	4.66	0.552	Extremely satisfied
6. Suitable and easy understandable language	4.74	0.484	Extremely satisfied
7. Suitable and clear easy reading font and font size	4.67	0.567	Extremely satisfied
8. Interesting story that enhanced engagement	4.62	0.579	Extremely satisfied
9. Suitable video length	4.56	0.605	Extremely satisfied
10. Video encouraged further learning	4.65	0.555	Extremely satisfied
11. The content was practical in daily life	4.78	0.418	Extremely satisfied
12. Overall satisfaction	4.68	0.429	Extremely satisfied

Satisfaction score: 5=extremely satisfied, 4=very satisfied, 3=moderately satisfied, 2=slightly satisfied, 1=not at all



Comparing awareness and knowledge of AMD before and after they underwent the self-learning AMD video, it was found that the mean score for awareness and knowledge of AMD of participants after they viewed the video ($\bar{x}=11.63$, S.D.=2.347) was significantly higher than the mean score for awareness and knowledge before viewing the video ($\bar{x}=3.78$, S.D.= 2.642) at $p<0.05$ ($t= 25.148$, P-value = 0.000). (Table 4) Results after viewing the self-learning L/Z role in eye health video also demonstrated the same trend. Mean score for awareness and knowledge of the role of L/Z in eye health ($\bar{x}=12.88$, S.D. = 9.474) was significantly higher than the mean score for awareness and knowledge before viewing the video ($\bar{x}=5.10$, S.D. =3.661) at $p<0.05$ ($t= 8.190$, P-value =0.000). (Table 5)

Table (4). Comparing the mean score for awareness and knowledge of AMD of participants before and after viewing the self-learning nutrition education video.

	before		after		T-test	Diff	P-value
	\bar{x}	S.D.	\bar{x}	S.D.			
Mean score for awareness and knowledge of AMD	3.78	2.642	11.63	2.347	-25.148***	102	0.000

*** statistically significant at .001

Table (5): Comparing the mean score for awareness and knowledge of the role of L/Z in eye health of participants before and after viewing the self-learning nutrition education video.

	before		after		T-test	Diff	P-value
	\bar{x}	S.D.	\bar{x}	S.D.			
Mean score for awareness and knowledge of the role of L/Z in eye health	5.10	3.661	12.88	9.474	-8.190***	102	0.000

*** statistically significant at .001

4. Discussion

According to the results, after viewing the AMD video, participants had the highest satisfaction score on the language followed by the content and the illustrations. They found the language demonstrated in the video was suitable and easy understandable, the content was practical in daily life and the illustration were clear and match to content. Satisfaction results also revealed the same trend for the L/Z role in eye health video, participants had the highest satisfaction score on the content followed by the illustrations



www.mecsj.com

and the language. They found the content which demonstrated in the video was practical in daily life, the illustrations can help to easily understand content and the language was suitable and easy understandable. This is concluded that the developed videos were well acceptable by the adults and elderly.

For the comparison between awareness and knowledge of AMD, and the role of L/Z in eye health before and after viewing videos, results revealed that the mean score for awareness and knowledge of AMD and the role of L/Z in eye health of participants were significantly higher than the mean score for awareness and knowledge before the videos ($p < 0.05$). Therefore, it could be inferred that the adults and elderly improved their awareness and knowledge of AMD and the role of L/Z in eye health after they underwent the self-learning intervention. The adults and elderly demonstrated a better understanding of AMD, the AMD risk factors, ways to detect AMD, the stages and treatment of AMD as well as the way to prevent AMD. Furthermore, the adults and elderly also learnt about the important role of nutrients in eye health, especially the role of L/Z in preventing AMD, the suggested daily intake for L/Z, rich source of L/Z, and a suggested meal plan that would provide adequate dietary L/Z. Results of this study were consistent with the study of Suthanya et al. (2010) which demonstrated that the knowledge of cataract patients and their caregivers were significantly different after receiving video education compared with their knowledge prior to receiving the education ($p < 0.01$). In addition, results of this study were in line with the study of Yanut et al. (2016) in individuals at risk of having stroke in northern Thailand communities. Individuals viewed a short (10.45 minutes) educational video about stroke, narrated in the appropriate northern dialect, most of them (80%) shown more knowledgeable about the risk factors and warning signs of strokes after viewing than prior to viewing the video.

5. Conclusion

The nutrition intervention has shown a positive impact on improving awareness and knowledge of AMD and the role of L/Z in eye health. Furthermore, the intervention was readily accepted by the adults and elderly, which enhanced their engagement and further learning. Health care practitioners, along with health promotion providers, especially elderly health promotions, may find this self-learning nutrition education as a tool which can be incorporated into their current education programs. This tool may be used in the hospital, health promotion centers, as well as health promotion units located in the community.



6. Acknowledgements

The study was supported by Panupong Wanjantuk, PhD, Faculty of Engineering, Khon Kaen University, Uruwan Yamborisut, PhD, Institute of Nutrition, Mahidol University and all the volunteers who participated in this research. The authors thank Ms. Susan Bowerman for the assistance of article edition. And finally, the authors would like to express our gratitude to all team members who assisted the study and the publication.

7. Conflicts of Interest

The authors declare no conflict of interest.

8. Reference

- Abdel-Aal E.M., H. Akhtar, K. Zaheer and R. Ali. (2013). Dietary sources of lutein and zeaxanthin carotenoids and their role in eye health. *Nutrients*, 5(4), 1169-1185.
- American Optometric Association. (2019). Lutein and Zeaxanthin - Eye-Friendly Nutrients. Retrieved from: <https://www.aoa.org/patients-and-public/caring-for-your-vision/nutrition/lutein-and-zeaxanthin>
- Busaba Tharasena & Siriporn Lawan. (2012). *Content of beta-Carotene, xanthophyll, lutein and zeaxanthin in vegetable as Thai side dish*. International Conference on Nutrition and Food Sciences; IPCBEE, IACSIT Press, Singapore. Retrieved from: <http://www.ipcbee.com/vol39/048-ICNFS2012-N3014.pdf>
- Food and Agriculture Organization of the United Nations. (2019). Nutrition education for the public. Discussion papers of the FAO Expert Consultation: Training needs for nutrition education: Guidelines for in-service training of nutrition educators Retrieved from: <http://www.fao.org/docrep/W3733E/w3733e05.htm>
- ForeseeHome, AMD Monitoring Program. (2019). Symptom of wet AMD. Retrieved from: <https://www.foreseehome.com/about-amd/>
- Health Systems Research Institute. (2019). Thai National Health Examination Survey (NHES V). Retrieved from: <http://kb.hsri.or.th/dspace/handle/11228/4604>



www.mecej.com

- Higgins MM, Barkley MC. (2003). Tailoring nutrition education intervention programs to meet needs and interests of older adults. *Journal of Nutrition for the Elderly*, 23(1): 59-79.
- Macula center. (2019). Cigarette and Macular degeneration. Retrieved from: <https://maculacenter.com/eye-news-tampa-bay/cigarettes-and-macular-degeneration/>
- Marse-Perlman J.A., A.I. Fisher, R. Klein, M. Palta, G. Block, A.E. Millen and J.D. Wright. (2001). Lutein and zeaxanthin in the diet and serum and their relation to age-related maculopathy in the third national health and nutrition examination survey. *American Journal of Epidemiology*, 153(5), 424-432.
- National Eye institute. (2019). National Eye Health Education Program (NEHEP). Retrieved from: https://nei.nih.gov/health/maculardegen/armd_facts
- OneBite Wellness. (2019). Eye Health Nutrition. Retrieved from: <https://onebitewellness.wordpress.com/2015/03/02/eye-health-nutrition/>
- Parker S, Powell L, Hermann J, Phelps J, Brown B. (2011). Preferred educational delivery strategies among limited income older adults enrolled in community nutrition education programs. *Journal of Extension*, 49(1), 1-10.
- Sahyoun NR, Pratt CA, Anderson A. Evaluation of nutrition education interventions for older adults: a proposed framework. (2004). *Journal of The American Dietetic Association*, 104(1), 58-69.
- Seddon J.M., U.A. Ajani, R.D. Sperduto, R. Hiller, N. Blair, T.C. Burton, M.D. Farber, E.S. Gragoudas et al. (1994). Dietary carotenoids, vitamins A, C, and E, and advanced age-related macular degeneration. Eye Disease Case-Control Study Group. *The Journal of the American Medical Association*, 272(18),1413-1420.
- Suthanya Nuanprasit, Pongpet Wuttipong and Kasira Juntraramanee. (2010). Effect of the video program of self-care knowledge and self-care ability of cataract patients and their caregivers. *Thai Journal of Nursing Council*, 25(2), 78-86.



www.mecsj.com

- Talley eye institute. (2019). Macular Hole. Retrieved from:
<https://talleyeyeinstitute.com/eye-conditions/macular-hole/>
- Thammasat Eye Center. (2019). Age-related Macular Degeneration. Retrieved from
<http://www.tec.in.th/index.php?page=TuEyeChannel>
- United States Department of Agriculture. (2017). Food Composition Database. Retrieved from: <https://ndb.nal.usda.gov/ndb/nutrients/index>
- Wantanee Yenjit, Prut Hanutsaha, Sapon Iamsirithaworn, Udomsiri Panrat, Pongsri Chusri and Chaweewan Yenjit. (2007). The fourth national survey of blindness, low vision and major eye diseases in Thailand. *Journal of Public Health Ophthalmology*, 21(1), 10-94.
- World Health Organization. Age-related Macular Degeneration Definition. (2019). Retrieved from: <https://www.who.int/blindness/causes/priority/en/index7.html>
- Yanat Wauters, Roshinee Oupra and Prakaikaew Tanasuwan. (2016). Development of a teaching model for prevention of stroke among people at risk in the northern Thai community. The Southern College Network. *Journal of Nursing and Public Health*, 3(2), 100-16.